

**DRAFT**

**INTERACTION PROFILE FOR:  
ATRAZINE, DEETHYLATRAZINE, DIAZINON, NITRATE, AND SIMAZINE**

Prepared by:

Syracuse Research Corporation  
Under Contract No. 205-97-0622

Prepared for:

**U.S. Department of Health and Human Services  
Public Health Service  
Agency for Toxic Substances and Disease Registry**

December, 2004

Comment Period Ends: March 31, 2005



## PREFACE

The Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) mandates that the Agency for Toxic Substances and Disease Registry (ATSDR) shall assess whether adequate information on health effects is available for the priority hazardous substances. Where such information is not available or under development, ATSDR shall, in cooperation with the National Toxicology Program, initiate a program of research to determine these health effects. The Act further directs that where feasible, ATSDR shall develop methods to determine the health effects of substances in combination with other substances with which they are commonly found. The Food Quality Protection Act (FQPA) of 1996 requires that factors to be considered in establishing, modifying, or revoking tolerances for pesticide chemical residues shall include the available information concerning the cumulative effects of substances that have a common mechanism of toxicity, and combined exposure levels to the substance and other related substances. The FQPA requires that the Administrator of the Environmental Protection Agency consult with the Secretary of the Department of Health and Human Services (which includes ATSDR) in implementing some of the provisions of the act.

To carry out these legislative mandates, ATSDR's Division of Toxicology (DT) has developed and coordinated a mixtures program that includes trend analysis to identify the mixtures most often found in environmental media, *in vivo* and *in vitro* toxicological testing of mixtures, quantitative modeling of joint action, and methodological development for assessment of joint toxicity. These efforts are interrelated. For example, the trend analysis suggests mixtures of concern for which assessments need to be conducted. If data are not available, further research is recommended. The data thus generated often contribute to the design, calibration or validation of the methodology. This pragmatic approach allows identification of pertinent issues and their resolution as well as enhancement of our understanding of the mechanisms of joint toxic action. All the information obtained is thus used to enhance existing or developing methods to assess the joint toxic action of environmental chemicals. Over a number of years, ATSDR scientists in collaboration with mixtures risk assessors and laboratory scientists have developed approaches for the assessment of the joint toxic action of chemical mixtures. As part of the mixtures program a series of documents, Interaction Profiles, are being developed for certain priority mixtures that are of special concern to ATSDR.

The purpose of an Interaction Profile is to evaluate data on the toxicology of the "whole" priority mixture (if available) and on the joint toxic action of the chemicals in the mixture in order to recommend approaches for the exposure-based assessment of the potential hazard to public health. Joint toxic action includes additivity and interactions. A weight-of-evidence approach is commonly used in these documents to evaluate the influence of interactions in the overall toxicity of the mixture. The weight-of-evidence evaluations are qualitative in nature, although ATSDR recognizes that observations of toxicological interactions depend greatly on exposure doses and that some interactions appear to have thresholds. Thus, the interactions are evaluated in a qualitative manner to provide a sense of what influence the interactions may have when they do occur.

The public comment period ends March 31, 2005. Comments should be sent to:

Agency for Toxic Substances and Disease Registry  
Division of Toxicology  
Attn: Hana Pohl, M.D., Ph.D.  
1600 Clifton Road, N.E.  
Mail Stop F-32  
Atlanta, GA 30333



## CONTRIBUTORS

### CHEMICAL MANAGER(S)/AUTHORS:

Hana Pohl, M.D., Ph.D.  
ATSDR, Division of Toxicology, Atlanta, GA

Joan Colman, Ph.D.  
Syracuse Research Corporation, Syracuse, NY



## PEER REVIEW

A peer review panel was assembled for this profile. The panel consisted of the following members:

Dale Hattis, Ph. D.  
The George Perkins Marsh Institute  
Center for Toxicology, Environment, and Development  
Clark University  
Worcester, MA

Kannan Krishnan, Ph. D.  
Department of Environmental and Occupational Health  
Faculty of Medicine  
University of Montreal, Montreal, QC, Canada

Sheldon Wagner, M.D.  
Department of Environmental and Molecular Toxicology  
Oregon State University  
Corvallis, OR

All reviewers were selected in conformity with the conditions for peer review specified in CERCLA Section 104(I)(13).

Scientists from ATSDR have reviewed the peer reviewers' comments and determined which comments will be included in the profile. A listing of the peer reviewers' comments not incorporated in the profile, with a brief explanation of the rationale for their exclusion, exists as part of the administrative record for this compound. A list of databases reviewed and a list of unpublished documents cited are also included in the administrative record.

The citation of the peer review panel should not be understood to imply its approval of the profile's final content. The responsibility for the content of this profile lies with the ATSDR.





## SUMMARY

Atrazine, deethylatrazine, simazine, diazinon, and nitrate were chosen as the subject mixture for this interaction profile because they frequently occur together in rural well water. Atrazine and simazine are triazine herbicides, deethylatrazine is a metabolite and an environmental degradation product of atrazine and other triazine herbicides, diazinon is an organophosphorus insecticide, and nitrate is a common contaminant resulting from fertilizers and human and animal waste. The exposures of greatest concern for this mixture in rural well water are intermediate and chronic oral exposures. No pertinent health effects data or physiologically-based pharmacokinetic (PBPK) models were located for the complete mixture. Therefore, the exposure-based screening assessment of potential health hazards for this mixture depends on an evaluation of the health effects and mechanistic data for the individual components and on the joint toxic action and mechanistic data for various combinations of the components. This profile discusses and evaluates the evidence for joint toxic action among atrazine, deethylatrazine, simazine, diazinon, and nitrate. The profile also discusses how public health assessments can incorporate concerns about interactions, additivity, and potential human exposures to mixtures of these chemicals.

Effects of concern for this mixture include reproductive effects (atrazine, deethylatrazine, and simazine), neurological effects (diazinon), and hematological effects (nitrate). Although none of the components has been classified as a carcinogen, atrazine and simazine can react with nitrite (nitrate metabolite) in the environment and *in vivo* to form N-nitrosoatrazine and N-nitrososimazine. Structure-activity considerations raise a concern for potential carcinogenicity of these nitrosamines.

To screen the mixture of atrazine, deethylatrazine, simazine, diazinon, and nitrate for potential hazards to public health, the hazard quotients (ratios of exposures to health guidance values) are estimated for the individual components. If only one or if none of the components has a hazard quotient that is at least 0.1, no further assessment of the *joint toxic action* is needed because additivity and/or interactions are unlikely to result in significant health hazard. If the hazard quotients for two or more of the mixture components equal or exceed 0.1, the following procedures are recommended. To screen this mixture for potential reproductive health hazard, an endpoint-specific hazard index for reproductive effects should be estimated for atrazine, deethylatrazine, and simazine (the triazine components of the mixture). The weight-of-evidence (WOE) analysis for interactions among these components indicates high confidence in the additivity assumption that is the basis for the hazard index. The potential effect of diazinon and nitrate on the reproductive toxicity of these triazines is uncertain. Separate hazard quotients are recommended to screen for the neurotoxicity of diazinon and the hematological toxicity of nitrate. The WOE analysis

indicates that because the triazine components may potentiate the neurologic toxicity of diazinon, the hazard quotient for diazinon may tend to underestimate the hazard of exposure to diazinon when these triazine components are present. Confidence in these predictions is medium. No information regarding the impact of interactions on the hematological toxicity of nitrate was available, so uncertainty regarding the impact of the other components on this effect of nitrate is high.

The interactions of atrazine and simazine with nitrite can result in the formation of N-nitrosoatrazine and N-nitrososimazine, which are more genotoxic than the parent triazine compounds. The WOE analysis predicts an increase in carcinogenic potential. Confidence in this prediction is medium; although adequate data are lacking for these specific compounds, data on other N-nitrosamines and their precursors support this conclusion. Further exposure-based screening for cancer risk is precluded by the lack of suitable data for N-nitrosoatrazine and N-nitrososimazine.

If the reproductive hazard index for the triazines or the hazard quotient for nitrate is greater than 1, or if the hazard quotient for diazinon is close to or above 1, then further evaluation is needed (ATSDR 2001a), using biomedical judgment and community-specific health outcome data. Community health concerns (ATSDR 1992) and the potential for carcinogenicity due to nitrosamine formation should be considered in further evaluations.

## TABLE OF CONTENTS

|  |      |
|--|------|
| PREFACE .....  | iii  |
| CONTRIBUTORS .....   | v    |
| PEER REVIEW .....  | vii  |
| SUMMARY .....  | ix   |
| TABLE OF CONTENTS .....  | xi   |
| LIST OF FIGURES .....  | xiii |
| LIST OF TABLES .....   | xiii |
| LIST OF ACRONYMS, ABBREVIATIONS, AND SYMBOLS .....   | xv   |
| 1. Introduction .....  | 1    |
| 2. Joint Toxic Action Data for the Mixture of Concern and Component Mixtures .....         | 5    |
| 2.1 Mixture of Concern .....   | 5    |
| 2.2 Component Mixtures .....   | 5    |
| 2.2.1 Atrazine/Deethylatrazine and Simazine .....  | 5    |
| 2.2.2 Atrazine and Diazinon .....  | 7    |
| 2.2.3 Simazine and Diazinon .....  | 9    |
| 2.2.4 Atrazine and Nitrate .....   | 9    |
| 2.2.5 Simazine and Nitrate .....   | 13   |
| 2.2.6 Diazinon and Nitrate .....   | 14   |
| 2.3 Relevance of the Joint Toxic Action Data and Approaches to Public Health .....         | 14   |
| 2.4 Recommendations for Data Needs .....   | 22   |
| 3. Recommendation for Exposure-Based Assessment of Joint Toxic Action of the Mixture ..... | 23   |
| 4. Conclusions .....   | 27   |
| 5. List of References .....  | 29   |
| Appendix A: Background Information for Atrazine and Deethylatrazine .....                  | 37   |
| A.1 Toxicokinetics .....   | 37   |
| A.2 Health Effects .....   | 38   |
| A.3 Mechanisms of Action .....   | 41   |
| A.4 Health Guidelines .....  | 42   |
| A.5 Derivation of Target Organ Toxicity Dose (TTD) Values .....                            | 44   |
| A.6 References .....   | 44   |
| Appendix B: Background Information for Simazine .....                                      | 47   |
| B.1 Toxicokinetics .....   | 47   |
| B.2 Health Effects .....   | 48   |
| B.3 Mechanisms of Action .....   | 48   |

|   |    |
|---|----|
| B.4 Health Guidelines .....   | 48 |
| B.5 Derivation of Target Organ Toxicity Dose (TTD) Values .....     | 49 |
| B.6 References .....  | 49 |
| Appendix C: Background Information for Diazinon .....               | 51 |
| C.1 Toxicokinetics .....  | 51 |
| C.2 Health Effects .....  | 51 |
| C.3 Mechanisms of Action .....                                      | 53 |
| C.4 Health Guidelines .....   | 54 |
| C.5 Derivation of Target-Organ Toxicity Dose (TTD) Values .....     | 54 |
| C.6 References .....  | 55 |
| Appendix D: Background Information for Nitrate .....                | 57 |
| D.1 Toxicokinetics .....  | 57 |
| D.2 Health Effects .....  | 58 |
| D.3 Mechanisms of Action .....                                      | 59 |
| D.4 Health Guidelines .....   | 60 |
| D.5 Derivation of Target Organ Toxicity Dose (TTD) Values .....     | 60 |
| D.6 References .....  | 61 |
| Appendix E: Chemical Structures of Organic Mixture Components ..... | 63 |

## LIST OF FIGURES

|  |    |
|--|----|
| Figure 1. Binary Weight-of-Evidence Scheme for the Assessment of Chemical Interactions . . . . . | 16 |
| Figure 2. Metabolic Pathways in Common to Atrazine and Simazine . . . . .                        | 39 |

## LIST OF TABLES

|  |    |
|--|----|
| Table 1. Effect of <b>Atrazine/Deethylatrazine</b> on <b>Simazine</b> : Reproductive Toxicity Effect of <b>Simazine</b> on <b>Atrazine/Deethylatrazine</b> : Reproductive Toxicity . . . . . | 17 |
| Table 2. Effect of <b>Atrazine/Deethylatrazine</b> on <b>Diazinon</b> : Neurological Toxicity . . . . .  | 18 |
| Table 3. Effect of <b>Simazine</b> on <b>Diazinon</b> : Neurological Toxicity . . . . .  | 19 |
| Table 4. Effect of <b>Atrazine</b> on <b>Nitrate</b> : Carcinogenicity Effect of <b>Nitrate</b> on <b>Atrazine</b> : Carcinogenicity . . . . .   | 20 |
| Table 5. Effect of <b>Simazine</b> on <b>Nitrate</b> : Carcinogenicity Effect of <b>Nitrate</b> on <b>Simazine</b> : Carcinogenicity . . . . .   | 21 |
| Table 6. MRLs and TTDs for Intermediate and Chronic Oral Exposure to Chemicals of Concern . . . .  | 24 |
| Table 7. Matrix of BINWOE Determinations for Intermediate or Chronic Simultaneous Oral Exposure to Chemicals of Concern . . . . .  | 25 |



## LIST OF ACRONYMS, ABBREVIATIONS, AND SYMBOLS

|                  |  |         |   |
|------------------|--|---------|---|
| ATSDR            | Agency for Toxic Substances and Disease Registry                               | NRC     | Nuclear Regulatory Commission                         |
|                  |  | NTP     | National Toxicology Program                           |
| BINWOE           | binary weight-of-evidence  | PAD     | population adjusted dose                              |
| CAS              | Chemical Abstracts Service   | 2-PAM   | pralidoxime   |
| CERCLA           | Comprehensive Environmental Response, Compensation, and Recovery Act           | PBPK    | physiologically based pharmacokinetic                 |
| CHO              | Chinese hamster ovary  | PBPK/PD | physiologically-based pharmacokinetic/pharmacodynamic |
| DACT             | diaminochlorotriazine  | ppb     | parts per billion                                     |
| DT               | Division of Toxicology   | RfC     | reference concentration                               |
| EC <sub>50</sub> | median effective concentration (produces desired effect in 50% of population)  | RfD     | reference dose transaminase                           |
| EPA              | Environmental Protection Agency  | SMR     | standardized mortality ratio                          |
| FQPA             | Food Quality Protection Act  | TTD     | target-organ toxicity dose                            |
| GnRh             | gonadotropin releasing hormone   | µg      | microgram   |
| IARC             | International Agency for Research on Cancer                                    | µmole   | micromole   |
| IRIS             | Integrated Risk Information System   | U.S.    | United States   |
|                  |  | VOC     | volatile organic compound                             |
|                  |  | WOE     | weight-of-evidence                                    |
| kg               | kilogram   | >       | greater than  |
|                  |  | ≥       | greater than or equal to                              |
| L                | liter  | =       | equal to  |
| LC <sub>50</sub> | median lethal concentration (produces desired effect in 50% of the population) | <       | less than   |
|                  |  | ≤       | less than or equal to                                 |
| LH               | luteinizing hormone  |         |   |
| LOAEL            | lowest-observed-adverse-effect level   |         |   |
| MCL              | maximum contaminant level  |         |   |
| MCLG             | maximum contaminant level goal   |         |   |
| mg               | milligram  |         |   |
| mM               | millimole  |         |   |
| MRL              | Minimal Risk Level   |         |   |
| NADH             | nicotinamide adenine dinucleotide phosphate                                    |         |   |
| ng               | nanogram   |         |   |
| NOAEL            | no-observed-adverse-effect level   |         |   |